arXiv:quant-ph/0103061v1 12 Mar 2001

Spin squeezing in nonlinear spin coherent states

Xiaoguang Wang
Institute of Physics and Astronomy, Aarhus University, DK-8000, Denmark
(February 1, 2008)

We introduce the nonlinear spin coherent state via its ladder operator formalism and propose a type of nonlinear spin coherent state by the nonlinear time evolution of spin coherent states. By a new version of spectroscopic squeezing criteria we study the spin squeezing in both the spin coherent state and nonlinear spin coherent state. The results show that the spin coherent state is not squeezed in the x, y, and z directions, and the nonlinear spin coherent state may be squeezed in the x and y directions.

Keywords: Spin squeezing, nonlinear spin coherent states

PACS numbers: 42.50.Dv

I. INTRODUCTION

In nonlinear systems such as optical Kerr medium [1], squeezed states of the radiation field [2] have been extensively studied. The spin squeezed states are also studied in nonlinear systems [3] and proved to be useful to enhance spectroscopic resolution [4], *e.g.*, in atomic clocks. The generation of the spin squeezed states has been studied by many ways, such as by interaction of atoms with squeezed light [4–8], quantum nondemolition measurement of atomic spin states [9], and atomic collisional interactions [10].

The spin squeezed states can be characterized in many different ways [3,4,11]. In our study we employ the criteria of spin squeezing recently proposed by Sørensen *et al.* [12] as a new version of spectroscopic squeezing [4,13]. The squeezing parameter is defined as [12]

$$\xi^2_{\vec{n}_1} = \frac{2j(\Delta J_{\vec{n}_1})^2}{\langle J_{\vec{n}_2}\rangle^2 + \langle J_{\vec{n}_3}\rangle^2}, \tag{1}$$

where $J_{\vec{n}} = \vec{n}\cdot\vec{J}$, $\vec{n}_i (i = 1, 2, 3)$ are orthogonal unit vectors and $\vec{J}$ is the spin-j angular momentum operator. The states with $\xi^2_{\vec{n}} < 1$ are spin squeezed in the direction $\vec{n}$. We will show a interesting feature that the squeezing parameters $\xi^2_x = \xi^2_y = \xi^2_z = 1$ for spin coherent states(SCS) [14], which indicates that the SCS is not squeezed in the x, y and z directions.

In this paper we introduce the nonlinear SCS and consider the spin squeezing in it. Sec. II gives the definition of the nonlinear SCS via its ladder operator formalism and propose a type of nonlinear SCS by the time evolution of the SCS under nonlinear Hamiltonian. In Sec. III we first prove that the SCS exhibits no squeezing in the x, y, and z directions and then study the spin squeezing in the nonlinear SCS. A conclusion is given in Sec. IV.

II. NONLINEAR SCS

We work in a $(2j+1)$-dimensional angular momentum Hilbert space $\{|j,m\rangle; m = -j, ..., +j\}$. It is convenient to define a number operator $\mathcal{N} = J_z + j$, and the 'number states' $|n\rangle$ which satisfy

$$|n\rangle \equiv |j, -j+n\rangle, \mathcal{N}|n\rangle = n|n\rangle. \tag{2}$$

The SCS is given by [14],

$$|\eta\rangle = (1+|\eta|^2)^{-j} \sum_{n=0}^{2j} \binom{2j}{n}^{1/2} \eta^n |n\rangle, \tag{3}$$

where η is complex.

It is easy to check that the SCS satisfies the following equation

$$J_-|\eta\rangle = \eta(2j - \mathcal{N})|\eta\rangle. \tag{4}$$

where the operators $J_\pm = J_x \pm iJ_y$. This is a ladder operator formalism of the SCS.

By recalling the definition of the bosonic nonlinear coherent state [15] and su(1,1) nonlinear coherent states [16], it is natural to define the nonlinear SCS as

$$f(\mathcal{N})J_-|\eta\rangle_{nl} = \eta(2j-\mathcal{N})|\eta\rangle_{nl}, \tag{5}$$

where $f(\mathcal{N})$ is a nonlinear function of the number operator $\mathcal{N}$. Eq.(5) is of the ladder operator form.

Now we propose a type of nonlinear SCS. The SCS under the evolution of the nonlinear Hamiltonian $F(\mathcal{N})$ is directly given by

$$\begin{aligned} |\eta, t\rangle &= e^{-itF(\mathcal{N})}|\eta\rangle \\ &= (1+|\eta|^2)^{-j} \sum_{n=0}^{2j} \binom{2j}{n}^{1/2} \eta^n e^{-itF(n)} |n\rangle. \end{aligned} \tag{6}$$

From Eqs.(4) and (6) we find the state $|\eta, t\rangle$ satisfies

$$e^{it[F(\mathcal{N}+1)-F(\mathcal{N})]} J_-|\eta, t\rangle = \eta(2j-\mathcal{N})|\eta, t\rangle. \tag{7}$$

According to the definition of the nonlinear SCS, the above state is a nonlinear SCS with the nonlinear function $e^{it[F(\mathcal{N}+1)-F(\mathcal{N})]}$.

For $F(\mathcal{N}) = \mathcal{N}^2 - \mathcal{N}$, Eq.(7) reduces to

$$e^{i2\mathcal{N}t}J_-|\eta,t\rangle=\eta(2j-\mathcal{N})|\eta,t\rangle. \tag{8}$$

Further let $t=\pi/2$, the above equation reduces to

$$\Pi J_-|\eta,\pi/2\rangle=\eta(2j-\mathcal{N})|\eta,\pi/2\rangle, \tag{9}$$

where $\Pi=(-1)^{\mathcal{N}}$ is the parity operator. Next we study the spin squeezing of the SCS $|\eta\rangle$ and the nonlinear SCS $|\eta,t\rangle$.

III. SPIN SQUEEZING

We first give a proof that there is no spin squeezing in the SCS along the x, y, and z directions, and then study the spin squeezing in the nonlinear SCS.

A. The SCS

In order to calculate the squeezing parameter (1) we need to know the expectation values $\langle\mathcal{N}^k\rangle$ and $\langle J_-^k\rangle(k=1,2)$. It is convenient to calculate $\langle\mathcal{N}^k\rangle$ by the generation function method. The generation function of the SCS is given by

$$G(\lambda)=\langle\eta|\lambda^{\mathcal{N}}|\eta\rangle=\frac{(1+\lambda|\eta|^2)^{2j}}{(1+|\eta|^2)^{2j}}, \tag{10}$$

from which the factorial moments follow

$$F(k)=\frac{d^kG(\lambda)}{d^k\lambda}|_{\lambda=1}=\frac{|\eta|^{2k}(2j)!}{(1+|\eta|^2)^k(2j-k)!}. \tag{11}$$

The factorial moments immediately give the expectation values of the operators $\mathcal{N}$ and $\mathcal{N}^2$ and the variance of $\mathcal{N}$

$$\langle\mathcal{N}\rangle=F(1)=\frac{2j|\eta|^2}{1+|\eta|^2}, \tag{12}$$

$$\langle\mathcal{N}^2\rangle=F(2)+F(1)=\frac{2j|\eta|^2+4j^2|\eta|^4}{(1+|\eta|^2)^2} \tag{13}$$

$$(\Delta\mathcal{N})^2=\frac{2j|\eta|^2}{(1+|\eta|^2)^2}. \tag{14}$$

From Eq.(3) the expectation value $\langle J_-^k\rangle$ are obtained as

$$\langle J_-^k\rangle=\frac{\eta^k(2j)!}{(1+|\eta|^2)^k(2j-k)!} \tag{15}$$

Now we calculate the squeezing parameter ξ_z^2, which is rewritten as

$$\xi_z^2=\frac{2j(\Delta\mathcal{N})^2}{|\langle J_-\rangle|^2}. \tag{16}$$

Then substituting Eqs.(14) and (15) into Eq.(16), we immediately obtain $\xi_z^2=1$.

To calculate ξ_x^2 and ξ_y^2 we need the identities

$$J_x^2=\frac{1}{4}[2j(2\mathcal{N}+1)-2\mathcal{N}^2+J_+^2+J_-^2], \tag{17}$$

$$J_y^2=\frac{1}{4}[2j(2\mathcal{N}+1)-2\mathcal{N}^2-J_+^2-J_-^2], \tag{18}$$

which gives the expectation values of J_x^2 and J_y^2 in terms of $\langle\mathcal{N}\rangle, \langle\mathcal{N}^2\rangle$ and $\langle J_\pm^2\rangle$.

From Eq.(15) and the relation $J_z=\mathcal{N}-j$, we get

$$\langle J_x\rangle=\frac{j(\eta+\eta^*)}{1+|\eta|^2},\ \langle J_y\rangle=\frac{j(\eta^*-\eta)}{i(1+|\eta|^2)},\ \langle J_z\rangle=\frac{j(|\eta|^2-1)}{1+|\eta|^2} \tag{19}$$

From Eqs.(12-13), (15), and (17-19), the variances of J_x and J_y are expressed as

$$\langle\Delta J_x\rangle^2=\frac{j(1+|\eta|^4-\eta^2-\eta^{*2})}{2(1+|\eta|^2)^2}=\frac{\langle J_y\rangle^2+\langle J_z\rangle^2}{2j}, \tag{20}$$

$$\langle\Delta J_y\rangle^2=\frac{j(1+|\eta|^4+\eta^2+\eta^{*2})}{2(1+|\eta|^2)^2}=\frac{\langle J_z\rangle^2+\langle J_x\rangle^2}{2j}. \tag{21}$$

The above two equations directly lead to $\xi_x^2=\xi_y^2=1$. Thus we have shown that the squeezing parameters $\xi_x^2=\xi_y^2=\xi_z^2=1$ for the SCS. That is to say, the SCS exhibits no squeezing in the x, y and z direction, irrespective of the complex η. We expect that the spin squeezing exists in the nonlinear SCS.

B. The nonlinear SCS

We examine the spin squeezing in the nonlinear SCS $|\eta,t\rangle$. The expectation values $\langle\ \mathcal{N}\rangle$,$\langle\mathcal{N}^2\rangle$ and the variance $(\Delta\mathcal{N})^2$ are time independent and given by Eqs.(12), (13), and (14), respectively.

From Eq.(6), we obtain the expectation value of J_-^k on the state $|\eta,t\rangle$ as

$$\langle J_-^k\rangle=\eta^k(1+|\eta|^2)^{-2j}\frac{(2j)!}{(2j-k)!}\times\sum_{n=0}^{2j-k}\binom{2j-k}{n}|\eta|^{2n}e^{it[F(n)-F(n+k)]}. \tag{22}$$

Of course Eq.(22) reduces to Eq.(15) at $t=0$.

By substituting Eqs.(14) and (22) into (16), the squeezing parameter ξ_z^2 is given by

$$\xi_z^2=\frac{1}{\left|\frac{1}{(1+|\eta|^2)^{2j-1}}\sum_{n=0}^{2j-1}\binom{2j-1}{n}|\eta|^{2n}e^{it[F(n)-F(n+1)]}\right|^2} \tag{23}$$

For $M+1$ complex quantities $c_i(i=0...M)$, there is an inequality

$$|c_0+c_1+...+c_M|\le|c_0|+|c_1|+...+|c_M| \tag{24}$$

Using this inequality in Eq.(23), we find that

$$\xi_z^2 \geq 1. \tag{25}$$

So a general conclusion is made that no squeezing occurs in the z direction for arbitrary nonlinear function $F(\mathcal{N})$. However spin squeezing may exist in the x or y directions. Next we make numerical calculations to show the spin squeezing.



FIG. 1. Squeezing parameters $\xi_\alpha^2(\alpha = x, y)$ as a function of time t. We plot ξ_α^2 for the nonlinear Hamtonian $F(\mathcal{N}) = \mathcal{N}^2$, $\xi_\alpha^3 + 0.6$ for $F(\mathcal{N}) = \mathcal{N}^3$, and $\xi_\alpha^2 + 1.2$ for $F(\mathcal{N}) = \mathcal{N}^4$. Solid line and dashed line correspond to ξ_x^2 and ξ_y^2, respectively. The parameters $\eta = 0.1$ and $j = 5$.

Fig.1 gives the squeezing parameters $\xi_\alpha^2(\alpha = x, y)$ as a function of time t for different nonlinear Hamiltonians $F(\mathcal{N}) = \mathcal{N}^k (k = 2, 3, 4)$. For small time t we observe that the state is squeezed in the x direction other than the y direction. As k increases, the frequency of occurrence of spin squeezing increases. In most of the time we also see that the spin squeezing alternatively appears in the x and y directions, i.e., when the state is squeezed in the $x(y)$ direction, it is not squeezed in the $y(x)$ direction. The state can show no spin squeezing in both the x and y directions in some small ranges of t, but it can not show spin squeezing at the same time in the two directions.



FIG. 2. Squeezing parameters $\xi_\alpha^2(\alpha = x, y)$ as a function of time t for the nonlinear Hamtonian $F(\mathcal{N}) = \sin(2\mathcal{N})$. Solid line and dashed line correspond to ξ_x^2 and ξ_y^2, respectively. We plot ξ_α^2 for $\eta = 0.1$, $\xi_\alpha^2 + 1$ for $\eta = 0.2$, and $\xi_\alpha^2 + 2.5$ for $\eta = 0.3$.The parameter $j = 5$.

It is interesting to consider the squeezing in the nonlinear Hamiltonian $H = \sin(a\mathcal{N})$ which can be realized in physical systems [17]. The numerical results are shown in Fig.2. For $\eta = 0.1$ we observe that the spin squeezing in the x and y directions appears alternatively in the beginning of the time evolution. For small time t, the state is squeezed in the y direction other than x direction in contrast to Fig.1. We also observe that the time range of spin squeezing decreases as the parameter η increases, i.e., the squeezing does not occur in most of the time as η is large.

IV. CONCLUSIONS

In conclusion we have given the definition and proposed an example of the nonlinear SCS. We have studied the spin squeezing in both the SCS and the nonlinear SCS. The main results are as follows:

1. The squeezing parameters $\xi_x^2 = \xi_y^2 = \xi_z^2 = 1$ for the SCS. That is to say, the SCS is not squeezed in the x, y and z directions, irrespective of the parameter η.
2. The nonlinear SCS shows no spin squeezing in the z direction for arbitrary nonlinear Hamiltonian $F(\mathcal{N})$.
3. The nonlinear SCS may be squeezed in the x and y directions. In most of the time the squeezing appears alternatively in the x and y directions as time goes on. Also we observe that the state can not be squeezed at the same time in the two directions.

The spin squeezing originates from the nonlinearity of the nonlinear SCS. Then we expect that the spin squeezing exists in other nonlinear SCS with difference nonlinear functions.

ACKNOWLEDGMENTS

The author thanks for many helpful discussions with Klaus Mølmer, Anders Sørensen, and Bin Shu. This work is supported by the Information Society Technologies Programme IST-1999-11053, EQUIP, action line 6-2-1.

---

[1] N. Imoto, H. A. Haus, and Y. Yamamoto, Phys. Rev. A **32**, 2287 (1985).

[2] D. F. Walls and G. J. Milburn, *Quantum Optics* (Springer, Berlin 1994).

[3] M. Kitagawa and M. Ueda, Phys. Rev. A **47**, 5138 (1993).

[4] D. J. Wineland *et al.*, Phys. Rev. A **46**, 11 (1992); Phys. Rev. A **50**, 67 (1994).

[5] G. S. Agarwal, R. R. Puri and R. P. Singh, Phys. Rev. A **56**, 2249 (1997).

[6] A. Kuzmich, K. Mølmer and E. S. Polzik, Phys. Rev. Lett. **79**, 4782 (1997); A. Kozhekin, K. Mølmer, and E. S. Polzik, Phys. Rev. A **62**, 033809 (2000).

[7] M. D. Lukin, S. F. Yelin, and M. Fleischhauer, Phys. Rev. Lett. **84**, 4232 (2000).

[8] L. Vernac, M. Pinard, and E. Giacobino, Phys. Rev. A **62**, 063812 (2000).

[9] A. Kuzmich, N. P. Bigelow, and L. Mandel, Europhys. Lett. **43**, 481 (1998); A. Kuzmich, L. Mandel, and N. P. Bigelow, Phys. Rev. Lett. **85**, 1594 (2000).

[10] A. Sørensen and K. Mølmer, Phys. Rev. Lett. **83**, 2274 (1999);

[11] D. A. Trifonov, J. Math. Phys. **34**, 100 (1993); **35**, 2297 (1994); Phys. Lett. A **187**, 284 (1994).

[12] A. Sørensen, L. M. Duan, J. I. Cirac and P. Zoller, Nature **409**, 63 (2001); L. M. Duan, A. Sørensen, J. I. Cirac and P. Zoller, Phys. Rev. Lett. **85**, 3991 (2000).

[13] G. S. Agarwal, Phys. Rev. A **49**, 4968 (1994).

[14] J. M. Radcliffe, J. Phys. A: Gen. Phys. **4**, 313 (1971); F. T. Arecchi, E. Courtens, R. Gilmore, and H. Thomas, Phys. Rev. A **6**, 2211 (1972); R. Gilmore, C. M. Bowden and L. M. Narducci, Phys. Rev. A **12**, 1019 (1975); L. M. Narducci, C. M. Bowden, V. Bluemel, G. P. Garrazana, and R. A. Tuft, Phys. Rev. A **11**, 973 (1975); G. S. Agarwal, Phys. Rev. A **24**, 2889(1981); K. Wòdkiewicz, Opt. Commun. **51**, 198 (1984); Phys. Rev. B **32**, 4750 (1985).

[15] R. L. de Matos Filho and W. Vogel, Phys. Rev. A **54**, 4560 (1996).

[16] X. Wang, Int. J. Mod. Phys. B. **14**, 1093 (2000); X Wang, B. C. Sanders, and S. H. Pan, J. Phys. A:Math.Gen. **33**, 7451 (2000).

[17] X. Wang, A. Sørensen and K. Mølmer, quant-ph/0012055.